



Renin-Angiotensin-Aldosterone System (RAAS)



Renin-angiotensin-aldosterone System (RAAS): The kidney begins to secrete renin, which catalyzes the conversion of angiotensinogen to angiotensin I, after blood pressure decreases. Subsequently, angiotensin I can be converted into angiotensin II under the catalytic action of angiotensin converting enzyme (ACE). Angiotensin II constricts blood vessels, leads to elevated blood pressure and stimulates the secretion of aldosterone, which has the effect of retaining sodium, discharging potassium and elevating blood pressure. Therefore, the detection of RAAS is of great significance for the diagnosis of hypertension.

Hypothalamic-pituitary-adrenal axis (HPA)



Hypothalamic-pituitary-adrenal axis (HPA): In the stress response of human body, the hypothalamus will release corticotropin releasing hormone (CRH) which stimulates the secretion of adrenocorticotrophic hormone (ACTH) from the pituitary. ACTH acts on the adrenal cortex to secrete cortisol. Meanwhile, cortisol has a feedback-inhibition to the hypothalamus and pituitary, achieving a self-stabilization. Adrenal diseases such as hypercortisolism, pheochromocytoma, primary hyperaldosteronism and congenital adrenal cortical hyperplasia can cause secondary hypertension and disorder of HPA axis. Therefore, the detection of HPA dysfunction is of great significance for the diagnosis of secondary hypertension.

Product	Pack Size	Cat.No.
Aldosterone	100 T	CMD0102
ACTH	100 T	CMD0202
Cortisol	100 T	CMD0302
Renin	100 T	CMD0402
Angiotensin II	100 T	CMD0502

INNOVATION · MANUFACTURE · INTEGRATION · SERVICE

AutoLumo Hypertension

Cat. No.: Aldosterone--CMD0102

ACTH--CMD0202

Cortisol--CMD0302

Renin--CMD0402

Angiotensin II--CMD0502

Autobio Diagnostics Co., Ltd.

offers more than 600 in vitro diagnostic products including CLIA (microplate based CLIA and magnetic particle based CLIA), ELISA, POCT (Point of Care Test), Microbiology and Biochemistry. As an ISO9001 and EN ISO13485 manufacturer, Autobio supplies high quality products through its well established sales network and is renowned as a reliable partner. For details please visit en.autobio.com.cn. Autobio Diagnostics Co., Ltd. | NO.87 Jingbei Yi Rd | National Eco&Tech Zone | Zheng zhou City | China 450016.

You can find us on:

Facebook: @AutobioGlobal

Youtube Channel: Autobio Diagnostics

Email: info@autobio-diagnostics.com

WhatsApp: +8618595767835

Tel: +86 371 6200 7036



AutoLumo Hypertension 20241209e

www.autobio.com.cn



Advantages

Identification of primary hypertension and secondary hypertension
Medication guidance for primary hypertension
Direct renin detection without sample pretreatment. Convenient and fast
Angiotensin II detection with a unique linking and amplification technique
Good correlation with international authoritative methods
Chemiluminescence analysis with high sensitivity, wide linear range



Renin

- ✘ Direct, quantitative detection of active renin in the specimens with monoclonal antibody
- ✘ Replace the traditional method of renin evaluation by calculating enzyme activity
- ✘ Trace to WHO international standard substance NTBSC68/356

Clinical Significance:

- 1 Great significance for the diagnosis of primary hyperaldosteronism
- 2 Diagnosis of renovascular hypertension due to renal artery stenosis
- 3 Provide effective information for the occurrence of cardiovascular complications in primary hypertension patients
- 4 Help clinicians interpret whether to conduct a renal vascular imaging

Performance Parameters:

- 1 Limit of Detection: 0.75 pg/mL
- 2 Linearity Range: 4 pg/ml - 500 pg/ml
- 3 Onboard Stability: 28 days
- 4 Repeatability: CV ≤ 10.0%

Note: Detailed parameters refer to the IFU of Renin CLIA Microparticles

Angiotensin II

- ✘ Convenient and fast test without sample extraction
- ✘ Unique linking and amplification technique

Clinical Significance:

- 1 Monitor the effects of drugs and evaluate medicine efficacy
- 2 Provide evidence for the classification and diagnosis of various kinds of hypertension and renal dysfunction
- 3 Auxiliary diagnosis of primary hyperaldosteronism, unilateral renal artery stenosis, adrenocortical hyperfunction, primary hypertension, etc.

Performance Parameters:

- 1 Limit of Detection: 4.00 pg/mL
- 2 Linearity Range: 10 pg/ml - 1000 pg/ml
- 3 Onboard Stability: 28 days
- 4 Repeatability: CV ≤ 10.0%

Note: Detailed parameters refer to the IFU of Angiotensin II CLIA Microparticles

Aldosterone

Clinical Significance:

- 1 Auxiliary diagnosis of primary hyperaldosteronism and secondary hyperaldosteronism
- 2 Auxiliary diagnosis of adrenocortical hypoactivities such as Addison's disease
- 3 Identification of selective hypoaldosteronism and primary hypoaldosteronism

Performance Parameters:

- 1 Limit of Detection: 16.32 pg/mL
- 2 Linearity Range: 40 pg/ml - 310 pg/ml
- 3 Onboard Stability: 28 days
- 4 Repeatability: CV ≤ 10.0%

Note: Detailed parameters refer to the IFU of Aldosterone CLIA Microparticles



Adrenocorticotrophic hormone (ACTH)

Clinical Significance:

Identification of hypercortisolism and functional status evaluation of hypothalamic-pituitary-adrenal axis (HPA)

- 1 Auxiliary diagnosis of adrenocortical tumor
- 2 Therapy observation of operation, radiotherapy and medication
- 3 Auxiliary diagnosis of pituitary-dependent hypercortisolism
- 4 Auxiliary diagnosis of ectopic ACTH syndrome

Identification of adrenocortical insufficiency:

- 1 Auxiliary diagnosis of primary adrenocortical hypofunction and congenital adrenocortical hyperplasia
- 2 Auxiliary diagnosis of secondary adrenal insufficiency caused by hypothalamic or adenohypophysis dysfunction

Performance Parameters:



- 1 Limit of Detection: 2.52 pg/mL
- 2 Linearity Range: 1 pg/ml - 2000 pg/ml
- 3 Onboard Stability: 28 days
- 4 Repeatability: CV ≤ 10.0%

Note: Detailed parameters refer to the IFU of ACTH CLIA Microparticles

Cortisol

Clinical Significance:

- 1 Auxiliary diagnosis of functional adrenal disease and Cushing's syndrome
- 2 Identification of primary or secondary adrenocortical hypofunction, such as Addison's disease, adrenal tuberculosis and adrenalectomy
- 3 Auxiliary diagnosis of hypofunction of anterior pituitary gland
- 4 Direct functional evaluation of adrenal gland and indirect functional evaluation of hypophysis



Performance Parameters:

- 1 Limit of Detection: 0.20 µg/dL
- 2 Linearity Range: 0.1 µg/dL - 60 µg/dL
- 3 Onboard Stability: 28 days
- 4 Repeatability: CV ≤ 10.0%

Note: Detailed parameters refer to the IFU of Cortisol CLIA Microparticles